

Complex interaction of polarized light with three-dimensional opal-based photonic crystals: diffraction and transmission measurements

M. V. Rybin¹, A. V. Baryshev^{1,2}, M. Inoue², A. A. Kaplyanskii¹,
V. A. Kosobukin¹, M. F. Limonov¹, A. K. Samusev¹, and A. V. Sel'kin¹,

¹ Ioffe Physico-Technical Institute, 194021, St. Petersburg, Russia.

² Toyohashi University of Technology, Toyohashi, Aichi 441–8580, Japan.

We investigated the photonic bandgap structure of synthetic opals by combining optical polarization-resolved diffraction patterns and polarization- and angle-resolved transmission spectra. The diffraction patterns were observed as symmetrical sets of spots, each being a colored fingerprint of the photonic bandgap for a certain direction [1]. In transmission spectra, the measured energy positions of the deeps were found to be in a good agreement with the calculated data. We found strong anisotropy in intensity of both diffracted and transmitted light with different polarizations along special crystallographic directions. The polarization-resolved diffraction patterns and transmission spectra are discussed in terms of the two-band mixing formalism [2] for Bloch states in a photonic crystal taking into account the relative orientation of different crystallographic planes $\{hkl\}$ in the opal structure. The conclusion has been made which bridges optical spectroscopy of photonic crystals and optical spectroscopy of conventional bulk homogeneous materials.

The support of the EC-funded projects PHOREMOST (FP6/2003/IST/2-511616) is gratefully acknowledged.

- [1] A. V. Baryshev *et al.*, *Phys. Rev. B*, **70**, 113104 (2004).
- [2] D. A. Mazurenko *et al.*, *Phys. Rev. Lett.* **91**, 213903 (2003).